

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 928 793 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
14.07.1999 Bulletin 1999/28

(51) Int. Cl.⁶: **C07D 417/06**, C07D 277/56,
C07D 277/48, A61K 31/425

(21) Application number: **98124819.8**

(22) Date of filing: **30.12.1998**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **02.01.1998 EP 98100007**


(71) Applicant:
F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)

(72) Inventors:
• **Alig, Leo**
4303 Kaiseraugst (CH)
• **Edenhofer, Albrecht**
4125 Riehen (CH)
• **Hilpert, Kurt**
4114 Hofstetten (CH)
• **Weller, Thomas**
4051 Basel (CH)

(74) Representative:
Witte, Hubert, Dr. et al
124 Grenzacherstrasse
4070 Basel (CH)

(54) **Thiazole derivatives**


(57) Compounds of formula I



as well as pharmaceutically usable salts and esters thereof, wherein R¹, R² and R³ have the significance given in claim 1, inhibit the binding of adhesive proteins to the surface of different types of cell and accordingly influence cell-cell and cell-matrix interactions. They can be used in the form of pharmaceutical preparations in the control or prevention of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi.


EP 0 928 793 A1

[0002] The present invention is concerned especially with thiazole derivatives of formula I




wherein

R¹ is





R^2 is



R³ is hydrogen, alkyl, cycloalkyl, aryl, aralkyl or heteroaryl;

R⁵, R⁶, R⁷ and R⁸ are each independently hydrogen, alkyl or cycloalkyl or R⁷ and R⁸ together with the N atoms to which they are attached form a 5- to 8-membered heterocyclic ring which can be substituted with alkyl;

R^9 is



R^{10} is aryl, aralkyl, heterocyclyl or an α -amino acid bonded via the amino group, a to i are zero or whole positive numbers, whereby a and b are each independently zero to 4, d is zero or 1, with f being equal to zero when d is equal to zero, the sum of c, d and e is ≥ 1 and ≤ 4 , f and h are each independently zero or 1, with i being other than zero when h = 1, the sum of g, h and i is ≥ 2 and ≤ 5 ;

and their pharmaceutically usable salts and esters.

[0003] The compounds of formula I and their pharmaceutically usable salts and esters are novel and have valuable pharmacological properties. In particular, they inhibit the binding of adhesive proteins such as fibrinogen, vitronectin, von Willebrand factor, fibronectin, thrombospondin and osteopontin to the vitronectin receptors (such as e.g. $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, etc.) on the surface on different types of cell. The said compounds therefore influence cell-cell and cell-matrix interactions. They can be used as vitronectin receptor antagonists in the prophylaxis or treatment of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi.

[0004] Objects of the present invention are the compounds of formula I and their aforementioned salts and esters per se and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments which contain the said compounds, their salts or esters, the use of the said compounds, solvates and salts as medicaments, especially for the prophylaxis and/or therapy of illnesses, e.g. in the treatment or prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi, and the use of the said compounds and salts and/or esters for the production of medicaments for the treatment or prophylaxis of, for example, neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi.

[0005] In the present description the term "alkyl", alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, preferably a straight-chain or branched-chain alkyl group with 1-4 carbon atoms. Examples of straight-chain and branched-chain C_1 - C_8 alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, preferably methyl, ethyl, isopropyl and tert.butyl.


[0006] The term "cycloalkyl", alone or in combination, signifies a cycloalkyl ring with 3 to 8 carbon atoms, preferably a cycloalkyl ring with 5 to 7 carbon atoms. Examples of C_3 - C_8 cycloalkyl rings are cyclopropyl, methyl-cyclopropyl, dimethyl-cyclopropyl, cyclobutyl, methyl-cyclobutyl, cyclopentyl, methyl-cyclopentyl, cyclohexyl, methylcyclohexyl, dimethyl-cyclohexyl, cycloheptyl and cyclooctyl, preferably cyclopentyl.

[0007] The term "alkoxy", alone or in combination, signifies an alkyl ether group in which the term "alkyl" has the previously given significance. Examples are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec.butoxy and tert.butoxy, with methoxy and ethoxy being preferred.

[0008] The term "aryl", alone or in combination, signifies a phenyl or naphthyl group which optionally carries one or more substituents each independently selected from alkyl, alkoxy, halogen, hydroxy, amino and the like, such as phenyl, p-tolyl, 4-methoxyphenyl, 4-tert.butoxyphenyl, 4-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 4-hydroxyphenyl, 1-naphthyl and 2-naphthyl. Phenyls and chlorophenyls are preferred, especially phenyl and ortho-, meta- and para-monochlorophenyls.

[0009] The term "aralkyl", alone or in combination, signifies an alkyl or cycloalkyl group as previously defined in which one hydrogen atom is replaced by an aryl group as previously defined, such as benzyl, 2-phenylethyl and the like, with benzyl being preferred.

[0010] The term "Het" denotes the hydrocyclic system



in R^2 , i.e. a saturated, partially unsaturated or aromatic 5- to 8-membered heterocycle which carries R^9 as a substituent and which can contain in addition to the nitrogen atom an additional nitrogen, oxygen or sulphur atom, with the oxygen atom being preferred of these. If desired, it can be substituted on one or more carbon atoms by halogen, alkyl, cycloalkyl, alkoxy, oxo etc. and/or on a secondary nitrogen atom (i.e. -NH-) by alkyl, cycloalkyl, aralkoxycarbonyl, alkanoyl, phenyl or phenylalkyl. Examples of such heterocyclic ring systems are the pyrrolidine, piperidine, piperazine, morpholine, thiamorpholine, pyrrole, imidazole, pyrazole or hexahydropyrimidine ring. Preferred are 5- or 6-membered rings and of these especially the pyrrolidine, piperidine and morpholine rings.

[0011] The term "heterocyclyl", alone or in combination, signifies a saturated, partially unsaturated or aromatic 5- to 10-membered heterocycle which contains one or more hetero atoms selected from nitrogen, oxygen and sulphur. If desired, it can be substituted on one or more carbon atoms by halogen, alkyl, alkoxy, oxo etc. and/or on a secondary

nitrogen atom (i.e. -NH-) by alkyl, cycloalkyl, aralkoxycarbonyl, alkanoyl, phenyl or phenylalkyl. Examples of such heterocyclyl groups are pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiamorpholinyl, pyrrolyl, imidazolyl (e.g. imidazol-4-yl and 1-benzoyloxycarbonylimidazol-4-yl), pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, hexahydropyrimidinyl, furyl, thienyl, thiazolyl, oxazolyl, thiazolyl, indolyl (e.g. 2-indolyl), quinolyl (e.g. 2-quinolyl, 3-quinolyl and 1-oxido-2-quinolyl), isoquinolyl (e.g. 1-isoquinolyl and 3-isoquinolyl), tetrahydroquinolyl (e.g. 1,2,3,4-tetrahydro-2-quinolyl), 1,2,3,4-tetrahydroisoquinolyl (e.g. 1,2,3,4-tetrahydro-1-oxo-isoquinolyl) and quinoxaliny. Preferred are 5- or 6-membered rings and of these especially piperidyl and pyridyl.


[0012] The term "heteroaryl", alone or in combination, signifies the aromatic compounds which fall under the definition of "heterocyclyl", with 5- and 6-membered rings, especially pyridyl, being preferred.

[0013] The term "amino", alone or in combination, signifies a primary, secondary or tertiary amino group bonded via the nitrogen atom, with the secondary amino group carrying an alkyl or cycloalkyl substituent and the tertiary amino group carrying two similar or different alkyl or cycloalkyl substituents or the two substituents on the nitrogen together forming a ring, such as, for example, NH₂, methylamino, ethylamino, dimethylamino, diethylamino, methyl-ethylamino, pyrrolidin-yl or piperidino etc. Primary amino, dimethylamino and diethylamino are preferred.


[0014] The term "halogen" signifies fluorine, chlorine, bromine or iodine, preferably chlorine.

[0015] Examples of "α-amino acids" bonded via the amino group are α-amino acids having the L- or D-configuration, the carboxyl group of which can be optionally derivatised as an ester or amide. Examples of such α-amino acids are L-valine, L-phenylalanine, L-leucine, L-isoleucine, L-serine, L-threonine, 3-(1-naphthyl)-L-alanine, 3-(2-naphthyl)-L-alanine, N-isopropyl-glycine, β-cyclohexyl-L-alanine and L-proline. Preferred are alanine, valine, phenylalanine, leucine and β-cyclohexyl-alanine, especially valine.


[0016] In the nomenclature used in the present description the ring atoms of the thiazole ring are numbered as follows:



with substituent R¹ being bonded to position 2 and substituent R² being bonded to position 4 and substituent R³ being bonded to position 5:



or substituent R² being bonded to position 5 and substituent R³ being bonded to position 4 of the thiazole ring:



In the most preferred arrangement R² is situated in position 5 and R³ is situated in position 4 of the thiazole ring.

[0017] Examples of physiologically usable salts of the compounds of formula I are salts with physiologically compatible mineral acids such as hydrochloric acid, sulphuric acid or phosphoric acid; or with organic acids such as methanesulphonic acid, acetic acid, trifluoroacetic acid, citric acid, fumaric acid, maleic acid, tartaric acid, succinic acid or

salicylic acid. The compounds of formula I having a free carboxy group can also form salts with physiologically compatible bases. Examples of such salts are alkali metal, alkaline earth metal, ammonium and alkylammonium salts such as the Na, K, Ca or tetramethylammonium salt. The compounds of formula I can also exist in the form of zwitterions.

[0018] The invention expressly includes pharmaceutically suitable derivatives of the compounds of formula I. For example, the COOH groups in R² can be esterified. Examples of suitable esters are the alkyl and aralkyl esters. Preferred esters are the methyl, ethyl, propyl, butyl, benzyl and (R/S)-1-((isopropoxy-carbonyl)-oxy)-ethyl esters. The ethyl esters are especially preferred.

[0019] The compounds of formula I can also be solvated, e.g. hydrated. The hydration can be effected in the course of the manufacturing process or can take place as a consequence of hygroscopic properties of an initially anhydrous compound of formula I.

[0020] The compounds of formula I can contain several asymmetric centres and can exist in the form of optically pure enantiomers, mixtures of enantiomers, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

[0021] Preferred compounds of formula I are those in which substituent R² is bonded to position 5 and substituent R³ is bonded to position 4 of the thiazole ring.

[0022] Also preferred are compounds of formula I in which Het in R² is a 5- or 6-membered heterocyclic ring system. Likewise preferred are those compounds of formula I in which Het is a 5- or 6-membered heterocyclic ring system in which an oxygen atom is present in the ring.

[0023] A group of especially preferred compounds of formula I comprises those in which Het in R² is a pyrrolidine, piperidine or morpholine ring and especially in which Het in R² is a pyrrolidine ring substituted with R⁹ in the 2- or 3-position, a piperidine ring substituted with R⁹ in the 3- or 4-position or a morpholine ring substituted with R⁹ in the 2- or 3-position.

[0024] Furthermore, preferred compounds are those in which R³ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl in which the substituted phenyl carries one or more substituents selected from alkyl, alkoxy, halogen, hydroxy, nitro and amino.

[0025] Likewise preferred compounds are those in which R⁴ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl in which the substituted phenyl carries one or more substituents selected from alkyl, alkoxy, halogen, hydroxy and amino.

[0026] A further group of preferred compounds comprises those in which R⁵, R⁶, R⁷ and R⁸ are hydrogen or R⁵ and R⁶ are hydrogen and R⁷ and R⁸ together with the N atoms to which they are attached form a 5- or 6-membered heterocyclic ring which can be substituted by alkyl, especially an imidazolidine or hexahydropyrimidine ring. Compounds in which R⁵, R⁶, R⁷ and R⁸ are hydrogen are especially preferred.

[0027] Furthermore, there are preferred compounds in which R¹⁰ is phenyl, pyridyl, substituted pyridyl or substituted phenyl, with the substituted phenyl and the substituted pyridyl carrying one or more substituents selected from alkyl, alkoxy, halogen, hydroxy and amino. Compounds in which R¹⁰ is phenyl are especially preferred. Compounds in which R¹⁰ is phenyl and f is equal to 0 are particularly preferred.

[0028] A group of preferred compounds of formula I comprises those in which substituent R² is bonded to position 5 and substituent R³ is bonded to position 4 of the thiazole ring, Het in R² is a 5- or 6-membered heterocyclic ring system in which R⁹ is bonded to one of the ring carbon atoms and in which optionally additionally an oxygen atom can be present in the ring, R³ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl, with the substituted phenyl carrying one or more substituents selected from alkyl, cycloalkyl, alkoxy, halogen, hydroxy, nitro and amino, R⁴ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl, with the substituted phenyl carrying one or more substituents selected from alkyl, alkoxy, halogen, hydroxy and amino, R⁵, R⁶, R⁷ and R⁸ are hydrogen or R⁷ and R⁸ together with the N atoms to which they are attached form a 5- or 6-membered heterocyclic ring which can be substituted by alkyl, R¹⁰ is phenyl, pyridyl, substituted pyridyl or substituted phenyl, with the substituted phenyl and the substituted pyridyl carrying one or more substituents selected from alkyl, alkoxy, halogen, hydroxy and amino, a and b each independently are zero to 3, the sum of c, d and e is ≥ 1 and ≤ 3 and the sum of g, h and i is ≥ 2 and ≤ 4 .

[0029] A group of especially preferred compounds of formula I comprises those in which substituent R² is bonded to position 5 and substituent R³ is bonded to position 4 of the thiazole ring, Het in R² is a pyrrolidine ring substituted with R⁹ in the 2- or 3-position, a piperidine ring substituted with R⁹ in the 3- or 4-position or a morpholine ring substituted with R⁹ in the 2- or 3-position, R³ is hydrogen, alkyl or phenyl, R⁴ is hydrogen, alkyl or phenyl, R⁵, R⁶, R⁷ and R⁸ are hydrogen or R⁷ and R⁸ together with the N atoms to which they are attached form an imidazolidine or hexahydropyrimidine ring and especially preferably R⁵, R⁶, R⁷ and R⁸ are hydrogen, R¹⁰ is phenyl and especially preferably R¹⁰ is phenyl and simultaneously f is equal to 0, a and b are each independently zero to 2 and the sum of g, h and i is equal to 2 or 3.

[0030] Examples of preferred compounds of formula I are:

Ethyl (RS)-3-[[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-propionate;

(RS)-3-[[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]propionic acid hydrochloride:

[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-4-yloxy]-acetic acid trifluoroacetate;

ethyl (RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]propionate;

(RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]-propionic acid;

(R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetic acid;

(S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetic acid;

ethyl (RS)-[1-(2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

(RS)-[1-(2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid hydrochloride;

ethyl (RS)- and (SR)-3-[[[(RS)-1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-3-phenyl-propionate];

(RS)- and (SR)-3-[[[(RS)-1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-3-phenyl-propionic acid];

ethyl rac [1-(2-guanidino-thiazole-4-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate hexafluorophosphate;

ethyl rac [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl rac [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl rac [1-(4-tert-butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl rac [1-(4-cyclopentyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

rac [1-(2-guanidino-thiazole-4-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(4-tert-butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(4-cyclopentyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

ethyl (S)-[1-(2-guanidino-4-methylthiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

ethyl rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-2-ylmethoxy]-acetate;

ethyl rac 3-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-propionate;

ethyl rac [4-(2-guanidino-4-methyl-thiazole-5-carbonyl)-morpholin-2-ylmethoxy]-acetate;

ethyl [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-4-ylmethoxy]-acetate;

(S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

(R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-2-ylmethoxy]-acetic acid;

rac 3-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-propionic acid;

rac [4-(2-guanidino-4-methyl-thiazole-5-carbonyl)-morpholin-2-ylmethoxy]-acetic acid;

[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-4-ylmethoxy]-acetic acid;

ethyl rac [1-(2-(3-benzyl-ureido)-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate;

rac [1-(2-(3-benzyl-ureido)-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

ethyl [1-(2-guanidino-thiazole-4-carbonyl)-piperidin-4-ylmethoxy]-acetate;

ethyl rac [1-(2-guanidino-4-methylthiazole-5-carbonyl)piperidin-3-yloxy]-acetate;

ethyl rac [1-[4-methyl-2-(3-methyl-ureido)-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]acetate;

[1-(2-guanidino-thiazole-4-carbonyl)-piperidin-4-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-yloxy]-acetic acid; and

rac [1-[4-methyl-2-(3-methyl-ureido)-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]-acetic acid.

[0031] Examples of especially preferred compounds of formula I are the following:

rac [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;


(R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;

rac [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid;


rac [1-(4-cyclopentyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid; and

rac [1-(4-tert-butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid.

[0032] Also an object of the invention is a process for the manufacture of a compound of formula I in which a thiazole-carboxylic acid of formula II




is reacted with an amine of formula III




or a salt thereof, with R^1 , R^3 , R^9 and Het having the previously given significance.

10 **[0033]** Especially preferred is a process in which the coupling of the thiazolecarboxylic acid with the amine component is effected by means of a coupling reagent and under the influence of a base. Especially suitable coupling reagents are, for example, benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) or O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU). Suitable bases are e.g. 4-methylmorpholine or N-methylmorpholine. All solvents which are inert under the given conditions can be used as the solvent. DMF is a preferred solvent for the reaction.

15 **[0034]** A further object of the invention comprises the intermediates of formula II



25 wherein R^1 and R^3 have the significances given above, with the proviso that, in the case of the acids, R^3 is not hydrogen or methyl when R^1 is



35 and, in the case of the esters, R^3 is not hydrogen, methyl or pyrid-4-yl N-oxide.

[0035] Particularly preferred intermediates are:

Ethyl 2-guanidino-4-propyl-thiazole-5-carboxylate hydrobromide;

40 ethyl 2-guanidino-4-phenyl-thiazole-5-carboxylate hydrobromide;

ethyl 4-tert-butyl-2-guanidino-thiazole-5-carboxylate hydrobromide;

ethyl 4-cyclopentyl-2-guanidino-thiazole-5-carboxylate hydrobromide;

45 2-guanidino-4-methyl-thiazole-5-carboxylic acid;

2-guanidino-4-propyl-thiazole-5-carboxylic acid hydrochloride;

50 2-guanidino-4-phenyl-thiazole-5-carboxylic acid;

4-tert-butyl-2-guanidino-thiazole-5-carboxylic acid hydrochloride;

4-cyclopentyl-2-guanidino-thiazole-5-carboxylic acid hydrochloride;

55 ethyl 2-(3-benzyl-ureido)-4-methyl-thiazole-5-carboxylate; .

2-(3-benzyl-ureido)-4-methyl-thiazole-5-carboxylic acid;

tert-butyl rac 3-carboxymethoxymethyl-piperidine-1-carboxylate;

tert-butyl (S)-3-carboxymethoxymethyl-piperidine-1-carboxylate;

5 tert-butyl (R)-3-carboxymethoxymethyl-piperidine-1-carboxylate;

tert-butyl rac 2-carboxymethoxymethyl-pyrrolidine-1-carboxylate;

ethyl rac (4-benzyl-morpholin-2-ylmethoxy)-acetate;

10

ethyl rac (morpholin-2-ylmethoxy)-acetate hydrochloride;

tert-butyl 4-ethoxycarbonylmethoxymethyl-piperidine-1-carboxylate;

15

ethyl (piperidin-4-ylmethoxy)-acetate;

tert-butyl rac 3-carboxymethoxy-piperidine-1-carboxylate;

ethyl rac (piperidin-3-yloxy)-acetate hydrochloride;

20

4-methyl-2-(3-methyl-ureido)-thiazole-5-carboxylic acid;

tert-butyl (RS)-3-(2-ethoxycarbonyl-ethylcarbamoyl)-piperidine-1-carboxylate;

25

ethyl (RS)-3-[(piperidine-3-carbonyl)-amino]-propionate hydrochloride;

tert-butyl (R)-[1-(R)-[1-phenyl-ethyl]-pyrrolidin-3-ylmethoxy]-acetate;

tert-butyl (R)-(pyrrolidin-3-ylmethoxy)-acetate;

30

tert-butyl (R)-[1-(2-guanidino-4-methylthiazole-5-carbonyl)pyrrolidin-3-ylmethoxy]-acetate;

tert-butyl (S)-[1-(R)-[1-phenyl-ethyl]-pyrrolidin-3-ylmethoxy]-acetate;

35

tert-butyl (S)-(pyrrolidin-3-ylmethoxy)-acetate;

tert-butyl (S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetate;

tert-butyl 3-(2-ethoxycarbonyl-1-phenyl-ethylcarbamoyl)-piperidine-1-carboxylate;

40

ethyl (RS)-3-phenyl-3-[(RS)-(piperidine-3-carbonyl)-amino]-propionate hydrochloride; and

tert-butyl [1-(2-guanidino-4-methylthiazole-5-carbonyl)-piperidin-4-yloxy]-acetate;

45

[0036] Compounds of formula I described above for use as therapeutically active substances are a further object of the invention.

[0037] Also an object of the invention are compounds of formula I described above for the production of medicaments for the prophylaxis and therapy of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptors.

50

[0038] Likewise an object of the invention are medicaments or pharmaceutical compositions containing a compound of formula I described above and a therapeutically inert carrier.

[0039] Also objects of the invention are the above medicaments which additionally contain one or more compounds selected from the group consisting of compounds of formula I, blood platelet inhibitors, anticoagulants, fibrinolytics as well as medicaments for the prophylaxis and therapy of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptors.

55

[0040] An object of the invention is also the use of the compounds of formula I described above for the production of medicaments e.g. for the treatment for prophylaxis of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptors.

[0041] Also an object of the invention is the use of one of the compounds of formula I described above for the production of medicaments for the treatment or prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi.

[0042] A further object of the invention comprises compounds of formula I when manufactured according to one of the described processes.


[0043] Likewise objects of the invention are methods for the treatment and prophylaxis of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptors and which comprise the administration of an effective amount of a compound of formula I.

[0044] An object of the invention is, further, a method for the treatment and prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi, whereby an effective amount of a compound of formula I described above is administered.


[0045] Also an object of the invention are compounds of formula I described above for the treatment and prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure and infections caused by viruses, bacteria or fungi.

[0046] The compounds of formula I hereinbefore and their pharmaceutically usable salts and esters are manufactured by the processes which are made available by the present invention. The substituents used in the following Schemes have the significance given above.

[0047] The compounds in accordance with the invention are manufactured by reacting a thiazolecarboxylic acid of the formula




with an amine of the formula



or a salt thereof with intermediary protection of the carboxy groups present in R⁹.

[0048] The coupling of the thiazolecarboxylic acid with the amine component is effected by means of a coupling reagent such as benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) or O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) under the influence of a base such as 4-ethylmorpholine or N-methylmorpholine (N-MM) in an inert solvent such as DMF at room temperature (Scheme 1).


Scheme 1



[0049] The subsequent liberation of the carboxy group protected as an ester is effected by means of a base such as aqueous LiOH or aqueous NaOH or also by cleavage using a strong acid such as concentrated hydrochloric acid or, in the case of a tert.butyl ester, trifluoroacetic acid.

[0050] For the preparation of the aforementioned thiazolecarboxylic acids (J. Med. Chem. 1991, 34, 914), conveniently an α -bromo-ketone of formula IV, such as e.g. a pyruvic acid ester, is reacted in a solvent such as ethanol with a thiourea derivative of formula V, such as 2-imino-4-thiobiuret, at elevated temperature (Scheme 2a).

Scheme 2a




[0051] A subsequent saponification of the ester group such as ethoxycarbonyl using a base such as aqueous NaOH or KOH yields a thiazole-4-carboxylic acid derivative of formula VI.

[0052] In another process variant, a substituted thiourea after cyclization to the thiazole can be reacted with an isocyanate such as benzyl isocyanate in a solvent such as DMF at room temperature, followed by a saponification of the ester as described above.

[0053] In a further process variant (Scheme 2b), which analogously to the process described above yields thiazole-5-carboxylic acid derivatives of formula in (Farmaco 1989, 44, 1011), there are used α -haloketones of formula VIII.

Scheme 2b

5




25 [0054] The α -haloketones of formula VIII are prepared from the corresponding β -ketoesters, such as ethyl butyrylacetate, ethyl pivaloylacetate, etc., by halogenation with e.g. bromine in a solvent such as water, conveniently at a temperature of 0-5°C (J. Chem. Soc. Perkin I 1982, 162).

30 [0055] For the preparation of the aforementioned amine components, an aminoalcohol protected with R⁰² in which R⁰² is a nitrogen protecting group such as Boc, benzyl or α -methylbenzyl is conveniently reacted with a bromoalkanoic acid ester, such as ethyl bromoacetate or ethyl bromopropionate, in the presence of a strong base such as aqueous NaOH (Scheme 3a).

Scheme 3a

35




[0056] A saponification of the ester function which may have taken place is counteracted either prior to the cleavage

or simultaneously with the cleavage of the nitrogen protecting group by esterification of the liberated carboxylic acid. The nitrogen protecting group Boc is cleaved off using hydrochloric acid in an alcohol equivalent to the ester, such as methanol or ethanol. The cleavage of the benzyl or α -methylbenzyl protecting group is accomplished by hydrogenation in ethanol in the presence of Pd/C. When a second nitrogen atom is present in the heterocycle, then both nitrogen atoms must each be bonded to different protecting groups.

[0057] In a process variant, the amine component can also be prepared from an aromatic precursor, such as e.g. pyridinylmethoxy-acetic acid, by hydrogenation in a solvent such as acetic acid and in the presence of Pd/C, preferably at elevated temperature and elevated pressure (Scheme 3b).


Scheme 3b



[0058] Esterification of the free carboxylic acid is effected according to known methods, for example using hydrochloric acid in an alcohol such as methanol or ethanol.

[0059] In a further process variant, an aminocarboxylic acid protected at the nitrogen with e.g. tertbutoxycarbonyl (Boc) or benzyloxycarbonyl (Cbz), such as e.g. N-Boc-piperidine-3-carboxylic acid, can be reacted with another amino-carboxylic acid protected as an ester, such as e.g. ethyl 3-amino-3-phenyl-propionate, or a salt thereof using a coupling reagent such as CDMT under the influence of a base such as N-MM in a solvent such as THF (Scheme 3c).

Scheme 3c



$R^{02} = \text{Boc, Cbz}$

$R^{03} = \text{alkyl, aralkyl}$

[0060] The nitrogen protecting group Boc is then cleaved off using hydrochloric acid in ethyl acetate and the protecting group benzyloxycarbonyl (Cbz) is cleaved off by hydrogenation in ethanol in the presence of Pd/C.

[0061] The conversion of a compound of formula I into a pharmaceutical usable salt can be carried by treating such a compound in the usual manner with an inorganic acid, for example a hydrohalic acid, such as, for example, hydrochloric acid or hydrobromic acid, sulphuric acid, nitric acid, phosphoric acid etc., or with an organic acid, such as, for example, acetic acid, citric acid, maleic acid, fumaric acid, tartaric acid, methanesulphonic acid or p-toluenesulphonic acid.

[0062] The corresponding carboxylate salts can be manufactured from the compounds of formula I by treatment with physiologically compatible bases.

[0063] The conversion of a compound of formula I into a pharmaceutically usable ester can be undertaken by esterifying such as compound in the usual manner or as described in the Examples.

[0064] As mentioned previously, the compounds of formula I and their pharmaceutically usable salts and esters inhibit especially the binding of various adhesive proteins such as fibrinogen, vitronectin, von Willebrand factor, fibronectin, thrombospondin and osteopontin to the vitronectin receptors (such as e.g. $\alpha v \beta 3$, $\alpha v \beta 5$, $\alpha v \beta 6$, $\alpha v \beta 8$, etc.) on the surface of various types of cell. The said compounds therefore influence cell-cell and cell-matrix interactions. Since the vitronectin receptors play a rôle, inter alia, in the spread of tumour cells, in the new growth of vascular tissue, in the degradation of bone tissue, in the migration of smooth muscle cells in vascular walls and in the penetration of virus particles into target cells, the said compounds can be used as vitronectin receptor antagonists in the control or prevention of neoplasms, tumor metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis fol-

lowing vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi. Since the binding of adhesive proteins to the fibrinogen receptor ($\alpha_{IIb}\beta_3$) on the surface of blood platelets is practically not inhibited, undesired side effects such as e.g. bleeding can be suppressed with the therapeutic application of the said compounds.

[0065] The inhibition of the binding of adhesive proteins such as e.g. fibrinogen to vitronectin receptors (such as e.g. $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, etc. or to the fibrinogen receptor ($\alpha_{IIb}\beta_3$) by compounds of formula I can be determined as described by L. Alig et al. (J.Med.Chem. 1992, 35, 4393-4407).

[0066] In detail thereto, the wells of microtitre plates (Nunc-Immunoplate MaxiSorp) were coated overnight at 4°C with the vitronectin receptor $\alpha_v\beta_3$ (from human placenta, 100 μ l/well) in a buffer system with 150 mmol/l NaCl, 1 mmol/l CaCl_2 , 1 mmol/l MgCl_2 , 0.0005% Triton X-100 and 20 mmol/l Tris HCl, pH 7.4. The non-specific binding sites were blocked by incubation with 3.5% bovine serum albumin (BSA from Fluka) at 20°C for at least 1 h. Before the beginning of the test the plates were washed in each case once with 150 mmol/l NaCl, 1 mmol/l CaCl_2 , 1 mmol/l MgCl_2 and 20 mmol/l Tris HCl, pH 7.4 (buffer A). The thus-coated plates can be stored for at least 2 months in the presence of 0.05% NaN_3 (in buffer A) at 4°C in a humidity chamber without loss of binding activity. Fibrinogen (IMCO, free from fibrinectin) was diluted to 1.5 μ g/ml in buffer A in the presence of 1% BSA. The wells coated with the receptor were incubated with fibrinogen (100 μ l/well) overnight at room temperature in the absence of or in the presence of increasing concentrations of RGDS (as the reference substance) or the compounds to be measured. Non-bound fibrinogen was removed by three-fold washing with buffer A and bound fibrinogen was detected by an ELISA procedure. Antibodies of rabbits directed against human fibrinogen (Dakopatts, Denmark), diluted in buffer A in the presence of 0.1% BSA, were added at room temperature for 1 h., followed by incubation with biotinylated antibodies directed against rabbit immunoglobulin (Amersham) for 30 min. Non-bound antibodies were removed by three-fold washing with buffer A. Thereafter, the pre-formed streptavidin-biotinylated peroxidase complex (Amersham) was added for 30 min. Three-fold washing with buffer A was again carried out. After addition of the peroxidase substrate ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid), Boehringer Mannheim) the enzyme activity was measured with a multichannel photometer (UVmax, Molecular Devices). The difference between total binding activity (in the absence of a test substance) and non-specific binding activity (in the presence of 100 μ M RGDS) is taken as the specific binding activity. The concentration of a test substance which is required to inhibit the specific binding activity by 50% was defined as the IC_{50} .

[0067] The isolation of the receptor $\alpha_v\beta_3$ used in the test can be carried out as follows: Human placenta is stored at -80°C immediately after its excision. In order to extract the receptor, each placenta is superficially thawed and cut into narrow strips with a scalpel. The pieces are washed twice with a buffer of 150 mmol/l NaCl, 1 mmol/l CaCl_2 , 1 mmol/l MgCl_2 and 20 mmol/l Tris HCl (pH 7.4). The proteins are extracted at room temperature for one hour with a buffer solution from 1% Triton X-100, 150 mmol/l NaCl, 1 mmol/l CaCl_2 , 1 mmol/l MgCl_2 , 20 mmol/l Tris HCl, 0.02% NaN_3 , 0.5 mmol/l phenylmethanesulphonyl fluoride, 1 mmol/l leupeptin and 2 mmol/l N-ethylmaleimide (pH 7.4) and filtered through sterile gauze. The filtrate is centrifuged at 30000 g for 30 min. at 4°C. The glycoproteins are firstly separated with the aid of a concanavalin A-Sepharose 4B column. The proteins bound to the column are eluted and then added to a Aeg-RGDS column. After repeated washing the bound vitronectin receptor is eluted by 3 mmol/l RGDS in a buffer of 0.1% Triton X-100, 150 mmol/l NaCl, 20 mmol/l Tris HCl, 1 mmol/l CaCl_2 , 1 mmol/l MgCl_2 , 0.05% NaN_3 (pH 7.0).

[0068] The results obtained in the foregoing test using representative compounds of formula as the test compound are compiled in the following Table.

Table 1

Substance	VNR
	IC_{50} [μ M]
rac [1-(2-Guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.018
(R)-[1-(2-Guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.05
rac [1-(4-Cyclopentyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.09
rac [1-(2-Guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.039
rac [1-(2-Guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.07
rac [1-(4-tert-Butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid	0.022

[0069] Preferred compounds have an IC_{50} value which is below 100 μ M; especially preferred compounds have a value below 10 μ M.

[0070] The compounds of formula I and their pharmaceutically usable salts and esters can be used as medicaments

(e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). The administration can, however, also be effected parentally such as intramuscularly or intravenously (e.g. in the form of injection solutions).

[0071] The compounds of formula I and their pharmaceutically usable salts and esters can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatine capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such adjuvants for tablets, dragées and hard gelatine capsules.

[0072] Suitable adjuvant for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

[0073] Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose, etc.

[0074] Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

[0075] Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

[0076] Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

[0077] In accordance with the invention the compounds of formula I and their pharmaceutical usable salts and esters can be used as vitronectin receptor antagonists, especially for the treatment or prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi. The dosage can vary in wide limits and will, of course be fitted to the individual requirements in each particular case. In the case of oral administration a daily dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to about 4 mg per kg body weight (e.g. approximately 300 mg per person), divided into preferably 1-3 individual doses, which can consist, for example, of the same amounts, should in general be adequate. It will, however, be clear that the upper limit given above can be exceeded when it is established that this is indicated.

[0078] The invention is illustrated in the following by Examples, with the Examples having no limiting character.

EXAMPLES

List of common abbreviations

[0079]

AcOEt	ethyl acetate
AcOH	acetic acid
Aeg-RGDS	aminoethylglycine-Arg-Gly-Asp-Ser-OH
Boc	tert-butoxycarbonyl
BOP	(benzotriazol-1-yloxy)-tris-(dimethylamino)-phosphonium hexafluorophosphate
BSA	bovine serum albumin
Cbz	benzyloxycarbonyl
CDMT	2-chloro-4,6-dimethoxy-1,3,5-triazine
DMF	dimethylformamide
EDC	N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide hydrochloride
EI	electron impact
ELISA	enzyme-linked immunosorbent assay
EtOH	ethanol
FAB	fast atom bombardment
HBTU	O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
ISP	ion spray (positively charged ions)
MeCN	acetonitrile
MeOH	methanol
MS	mass spectroscopy
N-MM	N-methylmorpholine
RGDS	H-Arg-Gly-Asp-Ser-OH
RP	reverse phase

RT	room temperature
m.p.	melting point
t-BuOH	tert-butanol
TFA	trifluoroacetic acid

5

Example 1

10 **[0080]** 13.9 ml of ethyl bromopyruvate are added to a solution of 11.81 g of 2-imino-4-thiobiuret (Aldrich) in 100 ml of ethanol and the reaction mixture is heated under reflux for 3 hours (J. Med. Chem. 34, 914-918 (1991)). Subsequently, the mixture is cooled to room temperature and the reaction product is precipitated by the addition of 550 ml of ethyl acetate and filtered off. There are obtained 14.6 g of yellowish ethyl 2-guanidino-thiazole-4-carboxylate hydrobromide. MS: 214 (M)⁺.

Example 2

15

[0081]

20 a) 1.62 ml of bromine are added dropwise within 10 minutes while stirring and cooling at 0-5°C to a 2-phase mixture of 5.06 ml of ethyl butyrylacetate and 14.4 ml of water (J. Med. Chem. 34, 914-918 (1991)). The mixture is stirred for a further 30 minutes at 0°C and then the product is extracted with ether. After drying there are obtained 7.6 g of crude bromoketone, which is used immediately in Example 3.

b) Analogously to the procedure of Example 2a, using ethyl benzoylacetate or ethyl pivaloylacetate or ethyl cyclopentylcarbonylacetate in place of ethyl butyrylacetate there is prepared the corresponding bromoketone.

25 Example 3

[0082] Analogously to the procedure of Example 1, using ethyl 2-chloroacetoacetate or the bromoketones prepared under Example 2 in place of ethyl bromopyruvate there are prepared the following compounds:

30 a) Ethyl 2-guanidino-4-methyl-thiazole-5-carboxylate hydrochloride, MS: 228 (M)⁺,

b) ethyl 2-guanidino-4-propyl-thiazole-5-carboxylate hydrobromide, MS: 256 (M)⁺,

35

c) ethyl 2-guanidino-4-phenyl-thiazole-5-carboxylate hydrobromide, MS: 290 (M)⁺,

d) ethyl 4-tert-butyl-2-guanidino-thiazole-5-carboxylate hydrobromide, MS: 271 (M+H)⁺,

40

e) ethyl 4-cyclopentyl-2-guanidino-thiazole-5-carboxylate hydrobromide, MS: 283 (M+H)⁺.

45 Example 4

[0083] 220 mg of 2-guanidino-4-methyl-thiazole-5-carboxylic acid, 265 mg of ethyl (RS)-3-[(piperidine-3-carbonyl)-amino]-propionate hydrochloride, 3 ml of DMF, 0.34 ml of N-methylmorpholine (N-MM) and 569 mg of O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) are stirred at RT for 22 hours. The reaction mixture is diluted with ethyl acetate and washed firstly with a dilute aqueous solution of sodium carbonate and sodium chloride, then with dilute sodium chloride solution and finally with saturated sodium chloride solution. The organic phase is dried over sodium sulphate acid evaporated in a vacuum. Chromatography on silica gel with ethyl acetate: ethanol 5:1 gives 270 mg of ethyl (RS)-3-[[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)piperidine-3-carbonyl]-amino]-propionate as a pale yellow foam. MS: 411 (M+H)⁺.

55 **[0084]** The starting materials can be prepared as follows:

a) 4.59 g of (RS)-piperidine-1,3-dicarboxylic acid 1-tert.-butyl ester, 3.51 g of 2-chloro-4,6-dimethoxy-(1,3,5)-triazine (CDMT), 60 ml of THF and 2.25 ml of N-MM are stirred under argon at 0°C for 3 hrs. After the addition of 3.07

g of β -alanine ethyl ester hydrochloride and 2.25 ml of N-MM the mixture is stirred at RT for 18 hrs. The reaction mixture is diluted with ethyl acetate and washed in sequence with ice-cold dilute hydrochloric acid, water, dilute sodium carbonate solution, water and saturated sodium chloride solution. After drying over sodium sulphate and evaporation of the solvent there are obtained 6.16 g of (RS)-3-(2-ethoxycarbonyl-ethylcarbamoyl)-piperidine-1-carboxylic acid tert.-butyl ester as a pale yellow oil. MS: 329 (M+H)⁺.

b) 985 mg of (RS)-3-(2-ethoxycarbonyl)-ethylcarbamoyl)-piperidine-1-carboxylic acid tert.-butyl ester are dissolved in 4.5 ml of ethyl acetate, treated with 4.5 ml of 4N HCl in ethyl acetate and stirred at RT for 1 hr. After evaporation of the solvent in a vacuum there are obtained 803 mg of ethyl (RS)-3-[(piperidine-3-carbonyl)-amino]-propionate hydrochloride (1:1), m.p. 105-108°C, MS: 299 (M+H)⁺.

c) 2.65 g of ethyl 2-guanidino-4-methyl-thiazole-5-carboxylate hydrochloride (prepared according to Example 3) are heated to 75°C for 7 hrs. in 70 ml of ethanol and 11 ml of 2N NaOH. The reaction mixture is evaporated to dryness in a vacuum. The residue is taken up in 40 ml of ethanol, the solution is filtered clear and the product is precipitated with 2 ml of acetic acid. There are obtained 1.82 g of 2-guanidino-4-methylthiazole-5-carboxylic acid of m.p. 196°C.

Example 5

[0085] 103 mg of ethyl (RS)-3-[[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-propionate are left to stand at RT in 2.1 ml of 25 percent hydrochloric acid for 6 hrs. The solution is evaporated and the residue is taken up in water and again evaporated. There are obtained 92 mg of (RS)-3-[[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-propionic acid hydrochloride (1:2) as a white foam, MS: 383 (M+H)⁺.

Example 6

[0086] 154 mg of tert.-butyl [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-4-yloxy]-acetate are dissolved in 1.5 ml of dichloromethane and treated with 1.5 ml of trifluoroacetic acid. After 2 hrs. the mixture is evaporated in a vacuum. The residue is dissolved in water and the solution is again evaporated. There are obtained 209 g of mg [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-4-yloxy]-acetic acid trifluoroacetate (1:2) as a light hygroscopic foam, MS: 342 (M+H)⁺.

[0087] The starting material (m.p. 205°C, MS: 398 (M+H)⁺) is obtained by coupling 2-guanidino-4-methyl-thiazole-5-carboxylic acid with tert.-butyl (piperidin-4-yloxy)-acetate according to the method given in Example 4.

Example 7

[0088] In analogy to Example 4, by coupling 2-guanidino-thiazole-4-carboxylic acid sodium salt with ethyl (RS)-3-[(piperidine-3-carbonyl)-amino]-propionate hydrochloride there is obtained ethyl (RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]-propionate as a white foam. MS: 397 (M+H)⁺.

Example 8

[0089] Analogously to Example 5, from ethyl (RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]-propionate there is obtained (RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]-propionic acid as the hydrochloride. This is neutralized in water with NH₃ and purified on Kieselgel 100 C₁₈-reverse phase. There is obtained (RS)-3-[[1-(2-guanidino-thiazole-4-carbonyl)-piperidine-3-carbonyl]-amino]-propionic acid as a white foam, MS: 369 (M+H)⁺.

Example 9

[0090] 140 mg of tert.-butyl (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetate are reacted as in Example 6. The crude trifluoroacetate is dissolved in water, neutralized with dilute NH₃, purified on Kieselgel 100 C₁₈-reverse phase and lyophilized from water. There are obtained 99 mg of (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetic acid, m.p. 136°C (sintering), [α]_D = +14.2°, (H₂O, c = 0.5), MS: 342 (M+H)⁺.

a) 20 mg of tetrabutylammonium bromide in 1 ml of water and then within 5 min. 20 g of 50% NaOH in water are added dropwise to a mixture of 2.05 g of (R)-1-[(R)- α -methylbenzyl]-3-pyrrolidinemethanol, 25 ml of toluene and 2.2 ml of tert.-butyl bromoacetate while stirring vigorously. After 6.5 hrs. the mixture is diluted with toluene, washed

neutral with water, dried and evaporated in a vacuum. Chromatography on silica gel gives 2.1 g of tert.-butyl (R)-[1-(R)-[1-phenyl-ethyl]-pyrrolidin-3-ylmethoxy]-acetate, $[\alpha]_D = +29.2^\circ$, (MeOH, $c = 1.0$), MS: 319 (M)⁺.

b) By catalytic hydrogenation on Pd/C in EtOH there is obtained therefrom tert.-butyl (R)-(pyrrolidin-3-ylmethoxy)-acetate, MS: 216 (M+H)⁺.

c) By coupling 2-guanidino-4-methyl-thiazole-5-carboxylic acid with tert.-butyl (R)-(pyrrolidin-3-ylmethoxy)-acetate according to the method given in Example 4 there is obtained tert.-butyl (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetate as a pale yellow resinous foam, $[\alpha]_D = +3.2^\circ$, (MeOH, $c = 0.5$), MS: 398 (M+H)⁺.

Example 10

[0091] Analogously to Example 9 there is prepared (S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetic acid; m.p. 139°C (sintering), $[\alpha]_D = -13.6^\circ$, (H₂O, $c = 0.5$), MS: 342 (M+H)⁺.

[0092] In addition, the following intermediates are prepared analogously:

a) tert.-Butyl (S)-[1-(R)-[1-phenyl-ethyl]-pyrrolidin-3-ylmethoxy]-acetate, $[\alpha]_D = +42.7^\circ$, (MeOH, $c = 1.0$), MS: 319 (M)⁺.

b) tert.-Butyl (S)-(pyrrolidin-3-ylmethoxy)-acetate, MS: 216 (M+H)⁺.

c) tert.-Butyl (S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-3-ylmethoxy]-acetate, pale yellow resinous foam, $[\alpha]_D = -2.4^\circ$, (MeOH, $c = 0.5$), MS: 398 (M+H)⁺.

Example 11

[0093] In the same manner as described in Example 4, from 2-guanidino-thiazole-5-carboxylic acid and ethyl (RS)-(piperidin-3-ylmethoxy)-acetate hydrochloride there is obtained ethyl (RS)-[1-(2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate, m.p. 156°C, MS: 370 (M+H)⁺.

[0094] 2-Guanidino-thiazole-5-carboxylic acid is obtained from ethyl 2-guanidino-thiazole-5-carboxylate by saponification with sodium hydroxide solution in alcohol, dilution with water and precipitation with hydrochloric acid at pH 3, m.p. 219°C.

Example 12

[0095] Analogously to Example 5, from ethyl (RS)-[1-(2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate after evaporation of the reaction solution and trituration of the residue in ether there is obtained (RS)-[1-(2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate hydrochloride (1:1), m.p. 75°C (dec.), MS: 342 (M+H)⁺.

Example 13

[0096] In analogy to Example 4, from 2-guanidino-4-methyl-thiazole-5-carboxylic acid and ethyl (RS)-3-phenyl-3-[(RS)-(piperidin-3-carbonyl)-amino]-propionate hydrochloride after chromatography on silica gel with dichloromethane:ethanol there is obtained a mixture of ethyl (RS)- and (SR)-3-[(RS)-1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-3-phenyl-propionate as a pale yellow foam, MS: 487 (M+H)⁺.

[0097] The starting material can be prepared as follows:

a) In analogy to Example 4a, from (RS)-piperidine-1,3-dicarboxylic acid 1-tert.-butyl ester and ethyl (RS)-3-amino-3-phenyl-propionate hydrochloride there is obtained tert.-butyl 3-(2-ethoxycarbonyl-1-phenyl-ethylcarbamoyl)-piperidine-1-carboxylate as a mixture of diastereomers, MS: 405 (M+H).

b) As given in Example 4b, there is obtained therefrom ethyl (RS)-3-phenyl-3-[(RS)-(piperidine-3-carbonyl)-amino]-propionate hydrochloride (1:1), MS: 304 (M)⁺.

Example 14

[0098] 243 mg of a mixture of ethyl (RS)- and (SR)-3-[(RS)-1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-

3-carbonyl]-amino]-3-phenyl-propionate are left to stand in 5 ml of 25% hydrochloric acid at RT for 24 hrs. The solution is evaporated, the residue is dissolved in water and the solution is adjusted to pH 8 with ammonia. The precipitate is filtered off under suction and purified by repeated trituration in water. There are obtained 77 g of a mixture of (RS)- and (SR)-3-[[[(RS)-1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidine-3-carbonyl]-amino]-3-phenyl-propionic acid, m.p. 174°C, MS: 459 (M+H)⁺.

Example 15

[0099] 146 ml of 3N sodium hydroxide solution are added to 14.6 g of the ester obtained under Example 1 and the reaction mixture is boiled under reflux for 3 hours (J. Med. Chem **34**, 914-918 (1991)). Then, the reaction mixture is cooled to RT, acidified with 73 ml of 6N hydrochloric acid and evaporated to ¼ of the volume. The precipitated material is filtered off and washed with water. After drying there are obtained 9.44 g of beige 2-guanidino-thiazole-4-carboxylic acid hydrochloride. MS: 186 (M)⁺.

Example 16

[0100] Analogously to the process in Example 15, from the esters obtained according to Example 3 there are prepared the following compounds:

a) 2-Guanidino-4-methyl-thiazole-5-carboxylic acid, MS: 200 (M⁺),

b) 2-guanidino-4-propyl-thiazole-5-carboxylic acid hydrochloride, MS: 229 (M+H)⁺,

c) 2-guanidino-4-phenyl-thiazole-5-carboxylic acid, MS: 263 (M+H)⁺,

d) 4-tert-butyl-2-guanidino-thiazole-5-carboxylic acid hydrochloride, MS: 243 (M+H)⁺,

e) 4-cyclopentyl-2-guanidino-thiazole-5-carboxylic acid hydrochloride, MS: 255 (M+H)⁺.

Example 17

[0101]

a) 4.05 ml of benzyl isocyanate are added to a solution of 5.0 g of ethyl 2-amino-4-methyl-thiazole-5-carboxylate in 50 ml of DMF. The reaction mixture is stirred at RT overnight, evaporated on a rotary evaporator and the residue is suspended in methylene chloride:methanol 1:1. The insoluble material is filtered off and dried. There are obtained 4.6 g of colourless ethyl 2-(3-benzyl-ureido)-4-methyl-thiazole-5-carboxylate, MS: 320 (M+H)⁺

b) A suspension of 3.6 g of the ester obtained under a) in 36 ml of ethanol is treated with 68 ml of 1N sodium hydroxide solution and boiled at reflux for 8 hours. Subsequently, the reaction mixture is poured into 70 ml of 1N ice-cold hydrochloric acid and the solution is evaporated to half of the volume. After cooling the separated crystals are filtered off and dried. There are thus obtained 2.25 g of colourless 2-(3-benzyl-ureido)-4-methyl-thiazole-5-carboxylic acid. MS: 292 (M+H)⁺.

Example 18

[0102] 200 ml of 50% sodium hydroxide solution and 1 g of butylammonium hydrogen sulphate are added to a solution of 21.5 g of tert-butyl rac 3-hydroxymethyl-piperidine-1-carboxylate (K. Hilpert et al., J. Med. Chem., 1994, 37, 3889; EP 0 468 231) in 200 ml of toluene. The 2-phase mixture is cooled to 15°C and treated with 30 ml of ethyl bromoacetate while stirring vigorously. After stirring at room temperature for 2.5 hours the reaction mixture is poured on to ice-water and extracted twice with ether. The organic phases are washed 4 times with water. The combined aqueous phases are acidified with concentrated hydrochloric acid and extracted twice with ethyl acetate. The ethyl acetate phases are washed with sodium chloride solution, dried and evaporated. There are obtained 18.3 g of tert-butyl rac-3-carboxymethyl-piperidine-1-carboxylate as a yellow oil, MS: 273 (M)⁺.

b) Hydrogen chloride is conducted for 10 minutes at 0°C into a solution of 18.3 g of the material obtained under a) in 183 ml of ethanol. Then, the reaction mixture is stirred at 0°C for a further 2 hours, then evaporated on a rotary evaporator and the residue is dried. There are obtained 9.6 g of pale beige crystalline ethyl rac (piperidin-3-ylmethoxy)-acetate hydrochloride.

MS: 202 (M+H)⁺.

Example 19

[0103]

a) The following compounds are prepared analogously to the procedure in Example 18a, but using a) tert-butyl (S)-3-hydroxymethyl-piperidine-1-carboxylate or b) tert-butyl (R)-3-hydroxymethyl-piperidine-1-carboxylate or c) rac-Boc-prolinol (EP 0 468 231) in place of tert-butyl rac 3-hydroxymethyl-piperidine-1-carboxylate:

a1) tert-Butyl (S)-3-carboxymethoxymethyl-piperidine-1-carboxylate,
MS: 273 (M)⁺,

b1) tert-butyl (R)-3-carboxymethoxymethyl-piperidine-1-carboxylate, MS: 273 (M)⁺,

c1) tert-butyl rac 2-carboxymethoxymethyl-pyrrolidine-1-carboxylate,
MS: 260 (M+H)⁺,

b) from the products a1, b1, c1 there are prepared analogously to the procedure in Example 18b the corresponding amino ester hydrochlorides a2), b2), c2), which are used immediately.

Example 20

[0104]

a) Analogously to the procedure described in Example 18a, but using ethyl bromopropionate in place of ethyl bromoacetate there is obtained tert-butyl rac 3-(2-ethoxycarbonyl-ethoxymethyl)-piperidine-1-carboxylate, MS: 316 (M+H)⁺,

b) Analogously to the procedure described in Example 18b), but using the products obtained according to Example 20a) there is obtained the corresponding free amine hydrochloride, which is used directly.

Example 21

[0105]

a) Analogously to the procedure described in Example 18a, but using rac 4-phenylmethyl-2-morpholine-methanol in place of tert-butyl 3-hydroxymethyl-piperidine-1-carboxylate there is obtained ethyl rac (4-benzyl-morpholin-2-ylmethoxy)-acetate.
MS: 294 (M+H)⁺

b) 1.0 g of the ester obtained under a) is dissolved in 10 ml of ethanol, treated with 3.4 ml of 1N hydrochloric acid and 0.1 g of palladium-on-charcoal and hydrogenated. After removal of the catalyst by filtration and evaporation of the filtrate there is obtained 0.8 g of ethyl rac (morpholin-2-ylmethoxy)-acetate hydrochloride, which is used immediately.

MS: 204 (M+H)⁺

Example 22

[0106]

a) A solution of 20.0 g of 4-pyridinylmethoxyacetic acid in 200 ml of acetic acid is treated with 2 g of platinum-on-charcoal and hydrogenated for 24 hours at 60°C under 100 bar of hydrogen. The catalyst is filtered off and the filtrate is evaporated. 24.5 g of residue are thus obtained.

b) Hydrochloric acid gas is conducted into a solution of the residue obtained under a) in 245 of ethanol at 0°C for 10 minutes. Then, the reaction mixture is stirred at 0°C for a further 2 hours, subsequently evaporated on a rotary evaporator and the residue is dried. 24.0 g of brown oil are obtained.

c) A solution of the product obtained under b) in 240 ml of dioxan is treated with 25.2 ml of 4-ethylmorpholine and a solution of 22.9 g of di-*t*-butyl dicarbonate in 50 ml of dioxan and stirred at RT overnight. The reaction mixture is evaporated, the residue is taken up in ethyl acetate and shaken once with 5% potassium hydrogen sulphate:10% potassium sulphate solution and twice with water. The organic phase is dried and evaporated and the residue is chromatographed on silica gel with hexane:ethyl acetate 4:1. 6.4 g of *tert*-butyl 4-ethoxycarbonylmethoxymethyl-piperidine-1-carboxylate are thus obtained.
MS: 301 (M)⁺.

d) The product obtained under c) is reacted analogously to the procedure described in Example 18b. There are thus obtained 5.4 g of ethyl (piperidin-4-ylmethoxy)-acetate,
MS: 202 (M+H)⁺.

Example 23

[0107] Analogously to Example 18, but using *tert*-butyl *rac* 3-hydroxy-piperidine-1-carboxylate in place of *tert*-butyl 3-hydroxymethyl-piperidine-1-carboxylate there are obtained the following two products:

a) *tert*-butyl *rac* 3-carboxymethoxy-piperidine-1-carboxylate, MS: 260 (M+H)⁺;

b) ethyl *rac* (piperidin-3-yloxy)-acetate hydrochloride, MS: 188 (M+H)⁺.

Example 24

[0108] 0.9 g of ethyl *rac* (piperidin-3-ylmethoxy)-acetate hydrochloride (Example 18b) is dissolved in 20 ml in dimethylformamide, treated with 2.18 ml of 4-ethylmorpholine, 1.0 g of the acid from Example 16e) and 1.52 g of benzotriazole-1-yloxy-tris(dimethyl-amino) phosphonium hexafluorophosphate (BOP). The reaction mixture is stirred at room temperature overnight and then evaporated on a rotary evaporator. The residue is chromatographed on a RP-18 column with a water:acetonitrile gradient. There is thus obtained 0.6 g of crystalline ethyl *rac* [1-(4-cyclopentyl-2-guanidino-thiazole-5-ylcarbonyl)-piperidin-3-ylmethoxy]-acetate.
MS: 438 (M+H)⁺.

Example 25

[0109] The following compounds were prepared analogously to Example 24 using the hydrochloride from Example 18 and the acids from Example 15 and Example 16:

a) Ethyl *rac* [1(2-guanidinothiazole-4-carbonyl)-piperidin-3-ylmethoxy]-acetate,
MS: 370 (M+H)⁺,

b) ethyl *rac* [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate hexafluorophosphate,
MS: 384 (M+H)⁺,

c) ethyl *rac* [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate, MS: 412 (M+H)⁺,

d) ethyl *rac* [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]acetate, MS: 446 (M+H)⁺,

e) ethyl *rac* [1-(4-*tert*-butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate, MS: 426 (M+H)⁺.

Example 26

[0110] 0.35 g of the ester obtained under Example 24 is suspended in 3.5 mol of tetrahydrofuran and treated with 2.4 ml of a 1N aqueous hydroxide solution. The mixture is stirred at room temperature for 2 hours, neutralized by the addition of 2.4 ml of 1N hydrochloric acid and evaporated on a rotary evaporator. After chromatography of the residue on a RP-18 column with a water:acetonitrile gradient there is obtained 0.32 g of colourless crystalline *rac* [1-(4-cyclopentyl-

2-guanidino-thiazol-5-ylcarbonyl)-piperidin-3-ylmethoxy]-acetic acid. MS: 410 (M+H)⁺.

Example 27

5 [0111] The following products are obtained analogously to Example 26, but using the esters from Example 25:

- a) rac [1-(2-Guanidino-thiazole-4-carbonyl)-piperidin-3-ylmethoxy]-acetic acid, MS: 342(M+H),
- b) rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid, MS: 356 (M+H)⁺,
- 10 c) rac [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid, MS: 384 (M+H)⁺,
- d) rac [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid, MS: 418(M+H)⁺,
- 15 e) rac [1-(4-tert-butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid, MS: 398 (M+H)⁺.

Example 28

[0112] The following compounds are prepared analogously to Example 24 using the hydrochlorides from Example 19
20 a2), b2), c2) and, respectively, Example 20 or Example 21b or Example 22d and the acid from Example 16a:

- a) Ethyl (S)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate,
MS: 384 (M+H)⁺,
- 25 b) ethyl (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetate,
MS: 384 (M+H)⁺,
- c) ethyl rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-2-ylmethoxy]-acetate,
MS: 370 (M+H)⁺,
- 30 d) ethyl rac 3-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-propionate,
MS: 398 (M+H)⁺,
- e) ethyl rac [4-(2-guanidino-4-methyl-thiazole-5-carbonyl)-morpholin-2-ylmethoxy]-acetate,
35 MS: 386 (M+H)⁺,
- f) ethyl [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-4-ylmethoxy]-acetate,
MS: 384 (M+H)⁺.

40 Example 29

[0113] The following products are obtained analogously to Example 26, but using the esters from Example 28:

- a) (S)-[1-(2-Guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
45 MS: 356 (M+H)⁺,
- b) (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
MS: 356 (M+H)⁺,
- 50 c) rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-pyrrolidin-2-ylmethoxy]-acetic acid,
MS: 342 (M+H)⁺,
- d) rac 3-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-propionsäure,
MS: 370 (M+H)⁺,
- 55 e) rac [4-(2-guanidino-4-methyl-thiazole-5-carbonyl)-morpholin-2-ylmethoxy]-acetic acid,
MS: 358 (M+H)⁺,

f) ethyl [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-4-ylmethoxy]-acetate,
MS: 356 (M+H)⁺.

Example 30

[0114] Analogously to Example 24, but using the amine from Example 18 b) and the acid from Example 17 b) in place of the acid from Example 15 there is obtained ethyl rac [1-[2-(3-benzyl-ureido)-4-methyl-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]-acetate, MS: 475 (M+H)⁺.

Example 31

[0115] The following acid is obtained analogously to Example 26, but using the ester from Example 30:

rac [1-[2-(3-Benzyl-ureido)-4-methyl-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]-acetic acid,
MS: 447 (M+H)⁺.

Example 32

[0116] Analogously to Example 24, but using the amine from Example 22d there is obtained

ethyl [1-(2-guanidino-thiazole-4-carbonyl)-piperidin-4-ylmethoxy]-acetate,
MS: 370 (M+H)⁺.

Example 33

[0117] Analogously to Example 24, but using the hydrochloride from Example 23 and the acid from Example 16a there is obtained

ethyl rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-yloxy]-acetate,
MS: 370 (M+H)⁺.

Example 34

[0118] Analogously to Example 24, but using the acid from Example 35 and the hydrochloride from Example 18b there is obtained

ethyl rac [1-[4-methyl-2-(3-methyl-ureido)-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]-acetate,
MS: 398 (M)⁺.

Example 35

[0119] Ethyl 4-methyl-2-[[[(methylamino)carbonyl]amino]-thiazole-5-carboxylate is hydrolyzed to the corresponding 4-methyl-2-(3-methyl-ureido)-thiazole-5-carboxylic acid, MS: 214 (M-H)⁻, analogously to Example 15.

Example 36

[0120] The following products are obtained analogously to Example 26, but using the ester from Example 32 or Example 33 or Example 34:

a) [1-(2-Guanidino-thiazole-4-carbonyl)-piperidin-4-ylmethoxy]-acetic acid,
MS: 342 (M+H)⁺,

b) rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-yloxy]-acetic acid,
MS: 342 (M+H)⁺,

c) rac [1-[4-methyl-2-(3-methyl-ureido)-thiazole-5-carbonyl]-piperidin-3-ylmethoxy]-acetic acid,
MS: 371 (M+H)⁺.

Example A

[0121] A compound of formula I can be used in a manner known per se as the active ingredient for the production of tablets of the following composition:

	Per tablet
Active ingredient	200 mg
Microcrystalline cellulose	155 mg
Corn starch	25 mg
Talc	25 mg
Hydroxypropylmethylcellulose	20 mg
	<u>425 mg</u>


Example B

[0122] A compound of formula I can be used in a manner per se as the active ingredient for the production of capsules of the following composition:

	Per capsule
Active ingredient	100.0 mg
Corn starch	20.0 mg
Lactose	95.0 mg
Talc	4.5 mg
Magnesium stearate	0.5 mg
	<u>220.0 mg</u>


Claims

1. Compounds of formula I




wherein

R¹ is



R² is





in which Het is a 5- to 8-membered heterocyclic ring system, which is substituted with R⁹ and in which an additional N, O or S atom can be present in the ring in addition to the nitrogen atom,

R³ is hydrogen, alkyl, cycloalkyl, aryl, aralkyl or heteroaryl;

R⁴ is hydrogen, alkyl, cycloalkyl, aryl or heteroaryl;

R⁵, R⁶, R⁷ and R⁸ are each independently hydrogen, alkyl or cycloalkyl or R⁷ and R⁸ together with the N atoms to which they are attached form a 5- to 8-membered heterocyclic ring which can be substituted with alkyl;

R⁹ is



R¹⁰ is aryl, aralkyl, heterocyclyl or an α -amino acid bonded via the amino group, a to i are zero or whole positive numbers, whereby a and b are each independently zero to 4, d is zero or 1, with f being equal to zero when d is equal to zero, the sum of c, d and e is ≥ 1 and ≤ 4 , f and h are each independently zero or 1, with i being other than zero when h = 1, the sum of g, h and i is ≥ 2 and ≤ 5 ; and their pharmaceutically usable salts and esters.

2. Compounds in accordance with claim 1, wherein R² is bonded to position 5 and R³ is bonded to position 4 of the thiazole ring.
3. Compounds in accordance with claim or 2, wherein Het in R² is a 5- or 6-membered heterocyclic ring system.
4. Compounds in accordance with any one of claims 1 to 3, wherein Het in R² is a pyrrolidine piperidine or morpholine ring.
5. Compounds in accordance with any one of claims 1 to 4, wherein R³ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl.
6. Compounds in accordance with any one of claims 1 to 5, wherein R⁴ is hydrogen, alkyl, cycloalkyl, phenyl or substituted phenyl.
7. Compounds in accordance with any one of claims 1 to 6, wherein R⁵, R⁶, R⁷ and R⁸ are hydrogen or R⁵ and R⁶


are each hydrogen and R^7 and R^8 together with the N atoms to which they are attached form a 5- or 6-membered heterocyclic ring.

8. Compounds according to any one of claims 1 to 7, wherein R^{10} is phenyl, pyridyl, substituted phenyl or substituted pyridyl.


9. Compounds according to claim 1 to 8, selected from:

rac [1-(4-tert-Butyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
 rac [1-(2-guanidino-4-propyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
 rac [1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
 (R)-[1-(2-guanidino-4-methyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid,
 rac [1-(2-guanidino-4-phenyl-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid and
 rac [1-(4-cyclopentyl-2-guanidino-thiazole-5-carbonyl)-piperidin-3-ylmethoxy]-acetic acid.

10. A process for the manufacture of a compound in accordance with any one of claims 1 to 9, which process comprises reacting a thiazolecarboxylic acid of formula II




with an amine of formula III




or a salt thereof, wherein R^1 , R^3 , R^9 and Het have the significance given in claim 1.

11. Compounds of formula II



wherein R^1 and R^3 are as defined in claims 1 to 8, and their salts and esters, with the proviso that, in the case of the acids, when R^1 is



R^3 is not hydrogen or methyl, and, in the case of the esters, R^3 is not hydrogen, methyl or pyrid-4-yl N-oxide.

12. Compounds in accordance with any one of claims 1-9 for use as therapeutically substances.

13. Compounds in accordance with any one of claims 1-9 for the production of medicaments for the prophylaxis and therapy of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptors.

5 14. A pharmaceutical preparation, containing a compound in accordance with any one of claims 1-9 and a therapeutically inert carrier.

10 15. A pharmaceutical preparation in accordance with claim 14, which additionally contains one or more compounds selected from the group consisting of compounds of formula I, blood platelet inhibitor's, anticoagulants, fibrinolytics and medicaments for the prophylaxis and therapy of illnesses which are caused by a malfunction of the binding of adhesive proteins to vitronectin receptor's.

16. The use of compound in accordance with any one of claims 1-9 for the production of medicaments.

15 17. The use of compounds in accordance with any one of claims 1-9 for the production of medicaments or the treatment and prophylaxis of neoplasms, tumour metastasing, osteoporosis, Paget's disease, diabetic retinopathy, macular degeneration, restenosis following vascular intervention, psoriasis, arthritis, kidney failure as well as infections caused by viruses, bacteria or fungi.

20 18. Compounds in accordance with any one of claims 1-9, when manufactured according to a process in accordance with claim 11.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 12 4819

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	US 4 649 146 A (T. TAKAYA ET AL.) 10 March 1987 * columns 5, 6, formulae Iq, X * * claims 1,8; examples 87,117 * ---	1,11,12, 14,16	C07D417/06 C07D277/56 C07D277/48 A61K31/425
A	EP 0 343 893 A (PFIZER INC.) 29 November 1989 * page 6, line 31 - line 40; claims 1,40; examples 12-14 * ---	1,12,14, 16,17	
A	DE 28 47 792 A (LEO PHARMACEUTICAL PRODUCTS LTD. A/S) 10 May 1979 * page 35, examples 4 and 5; page 38, examples 20 and 21 * page 12, line 4 - line 7; claims 1,17 * ---	1,12,14, 16,17	
A	PATENT ABSTRACTS OF JAPAN vol. 97, no. 4, 30 April 1997 & JP 08 337579 A (FUJISAWA PHARMACEUT CO LTD), 24 December 1996 ---	1,12,14, 16,17	
A	EP 0 417 751 A (FUJISAWA PHARMACEUTICAL CO., LTD.) 20 March 1991 * claims 1,16,18 * ---	1,12,14, 16	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07D
A	EP 0 050 458 A (PFIZER INC.) 28 April 1982 * claims 1,7 * ---	1,12,14, 16	
A	EP 0 161 841 A (PFIZER INC.) 21 November 1985 * claims 1,10 * ---	1,12,14, 16	
A	GB 2 138 803 A (SHIONOGI SEIYAKU K. K.) 31 October 1984 * page 1, Route B, formula (IV) * ---	11	
-/--			
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 30 April 1999	Examiner Hass, C
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 12 4819

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InLCI.6)
A,D	EP 0 468 231 A (F. HOFFMANN-LA ROCHE & CO. AG) 29 January 1992 -----		
			TECHNICAL FIELDS SEARCHED (InLCI.6)
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 30 April 1999	Examiner Hass, C
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P/MC01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 12 4819

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-04-1999

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4649146	A	10-03-1987	DK 41084 A	01-08-1984
			EP 0117082 A	29-08-1984
			JP 1868735 C	06-09-1994
			JP 59193878 A	02-11-1984
			US 4735957 A	05-04-1988

EP 343893	A	29-11-1989	AT 79114 T	15-08-1992
			AU 601905 B	20-09-1990
			AU 3509889 A	30-11-1989
			CA 1328871 A	26-04-1994
			CN 1037898 A,B	13-12-1989
			DD 283815 A	24-10-1990
			DK 249389 A	27-11-1989
			ES 2043012 T	16-12-1993
			FI 892498 A	25-11-1989
			GR 3005844 T	07-06-1993
			JP 2017181 A	22-01-1990
			JP 6078331 B	05-10-1994
			MX 16158 A,B	01-06-1993
			PT 90622 A,B	30-11-1989
			SU 1681728 A	30-09-1991
			US 4970318 A	13-11-1990
			YU 105989 A	31-08-1990

DE 2847792	A	10-05-1979	AT 366368 B	13-04-1982
			AT 773278 A	15-08-1981
			AU 517670 B	20-08-1981
			AU 4137678 A	17-05-1979
			BE 871807 A	07-05-1979
			CA 1120929 A	30-03-1982
			CH 645354 A	28-09-1984
			DK 493378 A,B,	08-05-1979
			FI 783327 A,B,	08-05-1979
			FR 2407930 A	01-06-1979
			GB 2009155 A,B	13-06-1979
			GR 65010 A	12-06-1980
			IE 47458 B	21-03-1984
			JP 54073784 A	13-06-1979
			LU 80476 A	15-06-1979
			NL 7811035 A	09-05-1979
			PH 16296 A	05-09-1983
			PT 68750 A	01-12-1978
			SE 446267 B	25-08-1986
			SE 7811457 A	08-05-1979
			US 4293549 A	06-10-1981
			ZA 7806175 A	31-10-1979

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 12 4819

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-04-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 417751 A	20-03-1991	AU 641425 B	23-09-1993
		AU 6242190 A	21-03-1991
		CA 2025356 A	16-03-1991
		CN 1050191 A	27-03-1991
		JP 1993240 C	22-11-1995
		JP 3141270 A	17-06-1991
		JP 7030056 B	05-04-1995
		PT 95317 A	22-05-1991
		US 5364871 A	15-11-1994
		US 5371097 A	06-12-1994
EP 50458 A	28-04-1982	US 4374843 A	22-02-1983
		AR 229703 A	31-10-1983
		AR 228674 A	30-03-1983
		AR 228675 A	30-03-1983
		AT 11048 T	15-01-1985
		AU 548992 B	09-01-1986
		AU 3206284 A	29-11-1984
		AU 530589 B	21-07-1983
		AU 7628781 A	22-04-1982
		CA 1156239 A	01-11-1983
		CS 235968 B	15-05-1985
		CS 235969 B	15-05-1985
		CS 235970 B	15-05-1985
		CS 8107500 A	13-02-1984
		DD 202293 A	07-09-1983
		DD 207718 A	14-03-1984
		DD 207719 A	14-03-1984
		DD 207913 A	21-03-1984
		DK 453181 A	15-04-1982
		FI 813177 A,B,	15-04-1982
		GR 75098 A	13-07-1984
		HK 52387 A	17-07-1987
		IE 51642 B	21-01-1987
		IN 163162 A	20-08-1988
		IN 163394 A	17-09-1988
		IN 163443 A	24-09-1988
		IN 163168 A	20-08-1988
		IN 163170 A	20-08-1988
		JP 1389402 C	23-07-1987
		JP 57095972 A	15-06-1982
		JP 61057307 B	06-12-1986
		KE 3707 A	03-04-1987
		PH 17562 A	01-10-1984
		PH 18467 A	18-07-1985
		PH 17856 A	09-01-1985

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 12 4819

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-04-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 50458 A		PT 73817 A,B	01-11-1981
		SU 1153829 A	30-04-1985
		SU 1311621 A	15-05-1987
		SU 1217258 A	07-03-1986
		SU 1205768 A	15-01-1983
		YU 105183 A	31-10-1983
		YU 105283 A	31-10-1983
		YU 105383 A	31-10-1983
		YU 246481 A	31-10-1983
		ZA 8107047 A	29-09-1982
		US 4452987 A	05-06-1984
		US 4510313 A	09-04-1985
		US 4590299 A	20-05-1986
EP 161841 A	21-11-1985	US 4560690 A	24-12-1985
		AT 44741 T	15-08-1989
		AU 554271 B	14-08-1986
		AU 4179085 A	07-11-1985
		CA 1262352 A	17-10-1989
		CS 248741 B	12-02-1987
		CS 248750 B	12-02-1987
		DD 233374 A	26-02-1986
		DK 190885 A,B,	31-10-1985
		EG 17391 A	30-12-1992
		FI 851683 A,B,	31-10-1985
		GR 851020 A	25-11-1985
		IN 165501 A	04-11-1989
		JP 1470507 C	14-12-1988
		JP 60239474 A	28-11-1985
		JP 63016387 B	08-04-1988
		MX 9203542 A,B	01-09-1992
		PH 21824 A	04-03-1988
		PT 80361 A,B	01-05-1985
		SU 1380614 A	07-03-1988
		SU 1400508 A	30-05-1988
		YU 72385 A	29-02-1988
		YU 114587 A	29-02-1988
GB 2138803 A	31-10-1984	NONE	
EP 468231 A	29-01-1992	AT 111890 T	15-10-1994
		AU 650458 B	23-06-1994
		AU 7949091 A	09-01-1992
		CA 2044636 A	06-01-1992
		DE 59103021 D	27-10-1994
		DK 468231 T	23-01-1995

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 12 4819

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-04-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 468231 A		ES 2061125 T	01-12-1994
		FI 913282 A	06-01-1992
		FI 961629 A	12-04-1996
		IE 66047 B	13-12-1995
		IL 98690 A	14-05-1996
		IL 112712 A	12-09-1996
		JP 1994338 C	22-11-1995
		JP 4230363 A	19-08-1992
		JP 7030022 B	05-04-1995
		NO 177704 B	31-07-1995
		NZ 238773 A	25-02-1994
		PT 98221 A,B	29-05-1992
		US 5393760 A	28-02-1995
		US 5532232 A	02-07-1996
		US 5595999 A	21-01-1997
		US 5583133 A	10-12-1996
		US 5763436 A	09-06-1998
		US 5260307 A	09-11-1993
		HR 930501 A	30-04-1996

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82